

On the Fairness of Additive Welfarist Rules*

Karen Frilya Celine
National University of Singapore
Singapore
karen.celine@u.nus.edu

Warut Suksompong
National University of Singapore
Singapore
warut@comp.nus.edu.sg

Sheung Man Yuen
National University of Singapore
Singapore
ysm@u.nus.edu

ABSTRACT

Background: Allocating indivisible goods is a ubiquitous task in fair division. We study *additive welfarist rules*, an important class of functions which choose an allocation that maximizes the sum of some function of the agents’ utilities. Prior work has shown that the maximum Nash welfare (MNW) rule is the unique additive welfarist rule that guarantees envy-freeness up to one good (EF1) [6, 21].

Objectives and Research Questions: If we restrict the class of instances—for example, to integer-valued instances—which additive welfarist rules guarantee EF1? Does MNW remain the only rule with this property, or are there other such rules? Note that integer-valued instances are used in practical applications such as Spliddit.

Methods: We consider several subclasses of instances including instances with identical goods, binary instances, two-value instances, normalized instances, and integer-valued instances. For each subclass, we attempt to characterize the additive welfarist rules that always yield EF1 allocations.

Results: We show that MNW remains the only additive welfarist rule that ensures EF1 for identical-good instances, two-value instances, as well as normalized instances with three or more agents, thus strengthening the earlier result by Suksompong [21]. On the other hand, if the agents’ utilities are integers, we demonstrate that several other rules offer the EF1 guarantee, and provide characterizations of these rules for various classes of instances.

Conclusions: Our results solidify the advantage of MNW over other additive welfarist rules for arbitrary real-valued instances. On the other hand, they indicate that when instances are integer-valued, the fairness provided by MNW is also offered by several other additive welfarist rules.

KEYWORDS

Welfarist rules; Fair division; Indivisible goods

ACM Reference Format:

Karen Frilya Celine, Warut Suksompong, and Sheung Man Yuen. 2026. On the Fairness of Additive Welfarist Rules. In *Appears at the 8th Games, Agents, and Incentives Workshop (GAIW-26). Held as part of the Workshops at the 25th International Conference on Autonomous Agents and Multiagent Systems., Paphos, Cyprus, May 2026*, IFAAMAS, 9 pages.

*This paper has been published at AAMAS 2025, and the full version (with all omitted proofs) can be found on arXiv [8].

Permission to make digital or hard copies of part or all of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for third-party components of this work must be honored. For all other uses, contact the owner/author(s).

Appears at the 8th Games, Agents, and Incentives Workshop (GAIW-26). Held as part of the Workshops at the 25th International Conference on Autonomous Agents and Multiagent Systems., Armstrong, Curry, Hosseini, Mattei, Tsang, Wqs (Chairs), May 2026, Paphos, Cyprus. © 2026 Copyright held by the owner/author(s).

1 INTRODUCTION

The distribution of limited resources to interested agents is a fundamental problem in society. Fairness often plays a key role in the distribution process—whether it be dividing inheritance between family members, allocating the national budget across competing sectors, or sharing credit and responsibility amongst participants in collaborative efforts. The complexity of this problem has led to the research area of *fair division*, which develops methods and algorithms to ensure that all agents feel fairly treated and has been studied in both economics and computer science [4, 18–20, 22].

What does it mean for an allocation of the resources to be “fair”? The answer depends on the context and the specific fairness benchmark applied. Among the plethora of fairness criteria that have been proposed, one of the most prominent is *envy-freeness*, which states that no agent should prefer another agent’s share to her own [11, 23]. In the ubiquitous setting of allocating *indivisible goods*—such as houses, cars, jewelry, and artwork—it is sometimes infeasible to attain envy-freeness. For example, if a single valuable good is to be allocated between two or more agents, only one of the agents can receive the good, thereby incurring envy from the remaining agents. In light of this, envy-freeness is often relaxed to *envy-freeness up to one good (EF1)*. The EF1 criterion allows an agent to envy another agent provided that the removal of some good in the latter agent’s bundle would eliminate the former agent’s envy. It is known that an EF1 allocation always exists, and that such an allocation can be computed in polynomial time [5, 6, 16].

In addition to fairness, another important property of allocations is *efficiency*. A well-known efficiency notion is *Pareto optimality (PO)*, which stipulates that no other allocation makes some agent better off and no agent worse off. In a far-reaching work, Caragiannis et al. [6] proved that a *maximum Nash welfare (MNW)* allocation, which maximizes the product of the agents’ utilities across all possible allocations,¹ satisfies both EF1 and PO, thereby offering fairness and efficiency simultaneously. In spite of this, the MNW rule does have limitations. For example, it avoids giving an agent zero utility at all costs—an allocation that gives most agents a tiny positive utility is preferred to another allocation that gives one agent zero utility and every other agent a large utility, as illustrated in the following example.

Example 1.1. Consider an instance with $n \geq 4$ agents and n goods g_1, \dots, g_n such that the utilities of the goods are as follows.

- $u_1(g_1) = n$.
- For $i \in \{2, \dots, n\}$, $u_i(g_{i-1}) = n - 1$ and $u_i(g_i) = 1$.
- $u_i(g) = 0$ for all other pairs (i, g) .

The only allocation \mathcal{A} that gives positive utility to every agent is the one that assigns good g_i to agent i for all $i \in \{1, \dots, n\}$ —this is also the MNW allocation. The sum of the agents’ utilities in this

¹This is equivalent to maximizing the sum of the logarithm of the agents’ utilities.

allocation is $n + (n - 1) = 2n - 1$. On the other hand, consider the allocation \mathcal{B} that gives good g_1 to agent 1, g_i to agent $i + 1$ for $i \in \{2, \dots, n - 1\}$, and g_n to agent n . The sum of the agents' utilities in this allocation is $n^2 - 2n + 3$, which is much larger than that in \mathcal{A} . Not only is \mathcal{B} also EF1 and PO, but it also ensures that one agent receives the same utility, all remaining agents except one receive much higher utility, while the exceptional agent receives only marginally lower utility.

The MNW rule belongs to the class of *additive welfarist rules*—rules that choose an allocation that maximizes the sum of some function of the agents' utilities for their bundles [18, p. 67]. Additive welfarist rules have the advantage that they satisfy PO by definition. Suksompong [21] showed that MNW is the only additive welfarist rule² that guarantees EF1 for all instances. Nevertheless, there exist other additive welfarist rules that guarantee EF1 for restricted classes of instances. For example, Montanari et al. [17, Appendix C] proved that the *maximum harmonic welfare (MHW)* rule ensures EF1 for the class of *integer-valued* instances. Note that this is an important class of instances in practice—for example, the popular fair division website Spliddit [12] only allows each user to specify integer values for goods summing up to 1000. In Example 1.1, allocation \mathcal{B} is the MHW allocation, and we saw earlier that \mathcal{B} has certain advantages compared to the MNW allocation \mathcal{A} . Beyond integer-valued instances, Eckart et al. [10] showed that certain additive welfarist rules based on *p-mean* functions, besides the MNW rule, also yield EF1 for the class of *normalized* instances with two agents—normalized instances have the property that the utility for the entire set of goods is the same for every agent. By the remark above, all instances on Spliddit are normalized.

In this paper, we study the necessary and sufficient conditions for additive welfarist rules to guarantee EF1 for different classes of instances. These classes include integer-valued instances, identical-good instances, two-value instances, normalized instances, as well as some of their combinations. We also establish relationships between the conditions for different classes of instances.

1.1 Our Results

In our model, we assume that each agent has an additive utility function over a set of indivisible goods. We consider additive welfarist rules, each of which is defined by a function f ; the rule chooses an allocation that maximizes the sum of the function f applied to the agents' utilities for their own bundles. In order for the additive welfarist rules to ensure that agents receive as much utility as possible, we assume that f is strictly increasing.³ For our characterizations, we only consider instances that are *positive-admitting*, i.e., instances in which there exists an allocation that gives positive utility to every agent.⁴ Our model is described formally in Section 2.

²Yuen and Suksompong [24] extended this result by showing that MNW is the only (not necessarily additive) welfarist rule that guarantees EF1 for all instances.

³If f is not strictly increasing, then the allocation chosen by the additive welfarist rule may not even be PO. In fact, we show in our full paper [8] that even if f is *non-decreasing* but not *strictly increasing*, then there exists an identical-good normalized instance such that the additive welfarist rule with function f does *not* always choose an EF1 allocation. This means that it is desirable for f to be strictly increasing.

⁴Even the MNW rule does not always return EF1 for instances that are not positive-admitting, unless there are additional tie-breaking mechanisms [6].

As one of the highlights of our paper, we extend the result of Suksompong [21] that the MNW rule is the unique additive welfarist rule that guarantees EF1 for all (positive-admitting) instances. In particular, we show that the same result still holds even if we restrict the class of instances to only identical-good instances, where each agent values every good equally but the value may differ between agents. This solidifies the advantage of MNW over other additive welfarist rules and indicates that the unique fairness of MNW stems from its *scale-invariance*,⁵ since for normalized identical-good instances, an additive welfarist rule with any strictly concave function f guarantees EF1. The proof of this result is technically involved, requiring a careful mathematical analysis of the growth rate of functions. The theorem statement can be found in Section 3.1.

To provide a more complete picture, we complement this result by studying other classes of instances. In Sections 3.2 and 3.3, we show that MNW remains the unique additive welfarist rule that ensures EF1 for the classes of two-value instances as well as normalized instances with three or more agents. Our result on normalized instances with three or more agents extends a result of Eckart et al. [10, Theorem 6], which only handles additive welfarist rules that are *p-mean* rules. For normalized instances with two agents, however, this characterization ceases to hold, and we provide a necessary condition and a sufficient condition for additive welfarist rules to guarantee EF1.

In Section 4, we consider integer-valued instances. For such instances, not only the MNW rule but also the MHW rule guarantees EF1 [17]. We shall attempt to characterize rules with this property. We show that for identical-good, binary, and two-value instances (which are also integer-valued), the respective functions defining the additive welfarist rules are characterized by Conditions 3, 4, and 5 respectively (see Definition 2.1). This allows much larger classes of functions beyond the MNW rule or the MHW rule. We provide examples and non-examples of *p-mean* rules as well as additive welfarist rules defined by modified logarithmic functions and modified harmonic functions that guarantee EF1 for these classes of instances. For the larger class of all integer-valued instances, we give a necessary condition and a sufficient condition for such additive welfarist rules.

In general, our characterizations are proven using the following outline: When a certain instance admits a unique EF1 allocation, an additive welfarist rule guaranteeing EF1 needs to return this allocation. In particular, moving some good to a different bundle in such a way that the allocation is no longer EF1 must decrease the social welfare. That is, the increase in welfare caused by adding the good to the new bundle must be strictly less than the decrease in welfare due to the removal of the good from the original bundle. This imposes a restriction on the growth rate of a function defining an additive welfarist rule guaranteeing EF1.

For all of our characterization results (Theorems 3.1, 3.5, 3.6, 4.1, 4.7, and 4.11), we show that if a function defining an additive welfarist rule guarantees EF1 for that particular class of instance for *some* number of agents n , then it also guarantees EF1 for that class for *every* number of agents n .

⁵That is, scaling the utility function of an agent by a constant factor does not change the outcome of the MNW rule.

Table 1: Conditions on the functions defining the additive welfarist rules that guarantee EF1 allocations for different classes of instances (see Definition 2.1). The logarithmic function corresponds to the MNW rule. The symbol “?” means that the precise condition remains unknown. For normalized instances with two agents, concaveness is a necessary condition (Proposition 3.7) and Condition 2 is a sufficient condition (Proposition 3.8). For integer-valued instances with no other restrictions, Condition 6a is a necessary condition (Proposition 4.14) and Condition 6b is a sufficient condition (Proposition 4.15).

	Identical-good	Binary	Two-value	Normalized	No other restrictions
Real-valued	Logarithmic (Th. 3.1)	Condition 4 (Th. 4.7)	Logarithmic (Th. 3.5)	$n \geq 3$: Logarithmic (Th. 3.6)	Logarithmic [6, 21]
Integer-valued	Condition 3 (Th. 4.1)	Condition 4 (Th. 4.7)	Condition 5 (Th. 4.11)	?	?




Figure 1: Relationships between the different conditions (see Definition 2.1).

Our results are summarized in Table 1. The relationships between the different conditions are summarized in Figure 1, and the relationships between additive welfarist rules that guarantee EF1 for different classes of instances are illustrated in Figure 2. All omitted proofs can be found in the full version of our paper [8].

2 PRELIMINARIES

Let $N = \{1, \dots, n\}$ be a set of $n \geq 2$ agents and $M = \{g_1, \dots, g_m\}$ be a set of goods. Each agent $i \in N$ has a *utility function* $u_i : 2^M \rightarrow \mathbb{R}_{\geq 0}$ such that $u_i(S)$ is i 's utility for a subset S of goods; we write $u_i(g)$ instead of $u_i(\{g\})$ for a single good $g \in M$. The utility function is *additive*, i.e., for each $S \subseteq M$, $u_i(S) = \sum_{g \in S} u_i(g)$; in particular, $u_i(\emptyset) = 0$. An *instance* consists of N , M , and $(u_i)_{i \in N}$.

An allocation $\mathcal{A} = (A_1, \dots, A_n)$ is an ordered partition of M into n bundles A_1, \dots, A_n such that A_i is allocated to agent $i \in N$. An allocation is *envy-free up to one good (EF1)* if for all $i, j \in N$ with $A_i \neq \emptyset$, there exists a good $g \in A_j$ such that $u_i(A_i) \geq u_i(A_j \setminus \{g\})$.

We say that an instance is

- *integer-valued* if $u_i(g) \in \mathbb{Z}_{\geq 0}$ for all $i \in N$ and $g \in M$;
- *identical-good* if for each $i \in N$, there exists $a_i \in \mathbb{R}_{>0}$ such that $u_i(g) = a_i$ for all $g \in M$;
- *binary* if $u_i(g) \in \{0, 1\}$ for all $i \in N$ and $g \in M$;
- *two-value* if there exist distinct $a_1, a_2 \in \mathbb{R}_{\geq 0}$ such that $u_i(g) \in \{a_1, a_2\}$ for all $i \in N$ and $g \in M$;⁶
- *normalized* if $u_i(M) = u_j(M)$ for all $i, j \in N$;
- *positive-admitting* if there exists an allocation (A_1, \dots, A_n) such that $u_i(A_i) > 0$ for all $i \in N$.

Let an instance be given. An *additive welfarist rule* is defined by a strictly increasing function $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R} \cup \{-\infty\}$ such that the rule chooses an allocation $\mathcal{A} = (A_1, \dots, A_n)$ that maximizes $\sum_{i \in N} f(u_i(A_i))$; if there are multiple such allocations, the rule chooses one arbitrarily. Since f is strictly increasing, $f(x) = -\infty$ implies that $x = 0$, which means that $f(x) \in \mathbb{R}$ for all $x \in \mathbb{R}_{>0}$.

⁶Two-value instances have received attention in fair division, particularly in relation to the MNW rule [1, 2].

A function f is *strictly concave* if for any distinct $x, y \in \mathbb{R}_{\geq 0}$ and $\alpha \in (0, 1)$, we have $f(\alpha x + (1 - \alpha)y) > \alpha f(x) + (1 - \alpha)f(y)$. We use \log to denote the natural logarithm, and define $\log 0$ as $-\infty$.

Definition 2.1. Let $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R} \cup \{-\infty\}$ be a strictly increasing function. For each $k \in \mathbb{Z}_{\geq 0}$, define $\Delta_{f,k} : \mathbb{R}_{>0} \rightarrow \mathbb{R} \cup \{\infty\}$ such that $\Delta_{f,k}(x) = f((k+1)x) - f(kx)$. The function f is said to satisfy

- *Condition 1* if $\Delta_{f,k}(b) > \Delta_{f,k+1}(a)$ for all $k \in \mathbb{Z}_{\geq 0}$ and $a, b \in \mathbb{R}_{>0}$;
- *Condition 1a* if $\Delta_{f,k}$ is a constant function (on domain $\mathbb{R}_{>0}$) for every $k \in \mathbb{Z}_{>0}$;
- *Condition 2* if $f(a) + f(b) < f(c) + f(d)$ for all $a, b, c, d \in \mathbb{R}_{\geq 0}$ such that $\min\{a, b\} \leq \min\{c, d\}$ and $ab < cd$;
- *Condition 3* if $\Delta_{f,k}(b) > \Delta_{f,k+1}(a)$ for all $k \in \mathbb{Z}_{\geq 0}$ and $a, b \in \mathbb{Z}_{>0}$;
- *Condition 3a* if $\Delta_{f,\ell}(b) > \Delta_{f,k}(a)$ for all $k, \ell \in \mathbb{Z}_{\geq 0}$ and $a, b \in \mathbb{Z}_{>0}$ such that $\ell < k$;
- *Condition 3b* if $\Delta_{f,k}(1) > \Delta_{f,k+1}(a) > \Delta_{f,k+2}(1)$ for all $k \in \mathbb{Z}_{\geq 0}$ and $a \in \mathbb{Z}_{>0}$;
- *Condition 4* if $\Delta_{f,k}(1) > \Delta_{f,k+1}(1)$ for all $k \in \mathbb{Z}_{\geq 0}$;
- *Condition 5* if $f((\ell+1)b + ra) - f(\ell b + ra) > \Delta_{f,k+1}(a) > \Delta_{f,k+2}(1)$ for all $a, b \in \mathbb{Z}_{>0}$ and $k, \ell, r \in \mathbb{Z}_{\geq 0}$ such that $a \geq b$ and $(k+1)b > \ell b + ra$;
- *Condition 6a* if $f((k+1)b - 1) - f(kb - 1) > \Delta_{f,k}(a)$ for all $k, a, b \in \mathbb{Z}_{>0}$;
- *Condition 6b* if $f(y + b) - f(y) > f(x + a) - f(x)$ for all $a, b \in \mathbb{Z}_{>0}$ and $x, y \in \mathbb{Z}_{\geq 0}$ such that $x/a \geq (y+1)/b$.

Note that for $k \in \mathbb{Z}_{\geq 0}$ and $x \in \mathbb{R}_{>0}$, we have $f((k+1)x) > f(kx)$ since f is strictly increasing, so $\Delta_{f,k}$ is well-defined and positive. Moreover, we have $\Delta_{f,k}(x) = \infty$ if and only if $k = 0$ and $f(0) = -\infty$.

Before proceeding further, we provide some intuition behind these conditions. Firstly, Condition 1a is a characterization of the logarithmic function $f(x) = \alpha \log x + \beta$ where $\alpha \in \mathbb{R}_{>0}$ and $\beta \in \mathbb{R}$: clearly, any logarithmic function satisfies Condition 1a, while for




Figure 2: Venn diagram showing the relationships between additive welfarist rules that guarantee EF1 for different classes of instances (see Table 1 for the full names of these classes). Each region, labeled by F with a subscript, represents the set of functions defining the additive welfarist rules that guarantee EF1 for the class of instances corresponding to the subscript. Some examples of these functions are given in the respective regions.

the converse, see Lemma 3.4. Most other conditions are related to Condition 1a in that they impose some restrictions on $\Delta_{f,k}$. For example, Conditions 3b and 5 allow $\Delta_{f,k}$ to take on any values as long as they lie within specific upper and lower bounds, whereas Condition 4 is equivalent to being concave on the integer domain. In other words, these conditions are satisfied by functions with properties similar to \log , with varying degrees of similarity depending on the conditions. Condition 2 is slightly different from the rest in that it references not only the logarithmic function (since $ab < cd$ is equivalent to $\log a + \log b < \log c + \log d$) but also the min function.

2.1 Examples of Additive Welfarist Rules

The *maximum Nash welfare (MNW) rule* is the additive welfarist rule defined by the function \log . We consider variations of this rule by defining, for each $c \in \mathbb{R}_{\geq 0}$, the *modified logarithmic function* $\lambda_c : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R} \cup \{-\infty\}$ such that $\lambda_c(x) = \log(x + c)$.

The *maximum harmonic welfare (MHW) rule* is the additive welfarist rule defined by the function $h_0(x) = \sum_{t=1}^x 1/t$. Note that this definition only makes sense when x is a non-negative integer; hence, we extend its definition to the non-negative real domain using the function $h_0(x) = \int_0^1 \frac{1-t^x}{1-t} dt$ [13]. We also consider variations of this rule. We define, for each $c \geq -1$, the *modified harmonic function* $h_c : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R} \cup \{-\infty\}$ such that⁷

$$h_{-1}(x) = \int_0^1 \frac{1-t^{x-1}}{1-t} dt \quad \text{and} \quad h_c(x) = \int_0^1 \frac{t^c - t^{x+c}}{1-t} dt$$

⁷This family of functions is defined for $c \in [-1, 0]$ in the domain of non-negative integers in previous work [17].

for $c > -1$. When the domain of h_c is restricted to the set of non-negative integers, we can rewrite⁸ the values as

$$h_{-1}(x) = \begin{cases} \sum_{t=1}^{x-1} \frac{1}{t} & \text{if } x \geq 1; \\ -\infty & \text{if } x = 0 \end{cases} \quad \text{and} \quad h_c(x) = \sum_{t=1}^x \frac{1}{t+c}$$

for $c > -1$ and all $x \in \mathbb{Z}_{\geq 0}$.

An *MNW allocation* (resp. *MHW allocation*) is an allocation chosen by the MNW rule (resp. MHW rule).

For $p \in \mathbb{R}$, the (*generalized*) *p-mean rule* is an additive welfarist rule defined by the function φ_p , where

$$\varphi_p(x) = \begin{cases} x^p & \text{if } p > 0; \\ \log x & \text{if } p = 0; \\ -x^p & \text{if } p < 0. \end{cases}$$

Hence, the MNW rule is also the 0-mean rule.

3 REAL-VALUED INSTANCES

We start with the general case where the utility of a good can be any (non-negative) real number. As mentioned earlier, the only additive welfarist rule that guarantees EF1 for all such instances is the MNW rule [21]. In this section, we consider the subclasses of *identical-good* instances (Section 3.1), *two-value* instances (Section 3.2), and *normalized* instances (Section 3.3). We demonstrate that the characterization of MNW continues to hold for the first two classes, as well as for the third class when there are three or more agents. To this end, we show that the only additive welfarist rule that ensures EF1 for these classes of instances is defined by the logarithmic function, which is equivalent to Conditions 1 and 1a.

3.1 Identical-Good Instances

Recall from Section 2 that identical-good instances have the property that each agent assigns the same utility to every good, although the utilities may be different for different agents. We state the characterization for such instances.

THEOREM 3.1. *Let $n \geq 2$ be given, and let $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R} \cup \{-\infty\}$ be a strictly increasing function continuous on $\mathbb{R}_{>0}$. Then, the following statements are equivalent:*

- For every positive-admitting identical-good instance with n agents, every allocation chosen by the additive welfarist rule with f is EF1.
- There exist constants $\alpha \in \mathbb{R}_{>0}$ and $\beta \in \mathbb{R}$ such that $f(x) = \alpha \log x + \beta$ for all $x \in \mathbb{R}_{\geq 0}$.

Note that the MNW rule is equivalent to the additive welfarist rule with a function that satisfies Theorem 3.1(b); therefore, the theorem essentially says that the only additive welfarist rule that guarantees EF1 for identical-good instances is the MNW rule.

The implication (b) \Rightarrow (a) follows from the result of Caragiannis et al. [6] that MNW guarantees EF1. We shall prove the implication (a) \Rightarrow (b) via a series of lemmas. We begin by showing that *Condition 1* is a necessary condition for a function f that satisfies the statement in Theorem 3.1(a).

⁸We prove this equivalence in the full version of our paper [8].

LEMMA 3.2. *Let $n \geq 2$ be given, and let $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R} \cup \{-\infty\}$ be a strictly increasing function such that the statement in Theorem 3.1(a) holds. Then, f satisfies Condition 1.*

PROOF. Let $k \in \mathbb{Z}_{\geq 0}$ and $a, b \in \mathbb{R}_{> 0}$ be given. Consider an instance with n agents and $(k+1)n$ goods such that the utility of each good is a for agent 1 and b for the remaining agents. This instance is an identical-good instance; moreover, it is positive-admitting since the allocation where every agent receives $k+1$ goods gives positive utility to every agent.

Let $\mathcal{A} = (A_1, \dots, A_n)$ be the allocation such that every agent receives $k+1$ goods. We have $\sum_{i \in N} f(u_i(A_i)) = f((k+1)a) + (n-1)f((k+1)b)$. Note that \mathcal{A} is the only EF1 allocation; indeed, if some agent does not receive exactly $k+1$ goods, then some agent i receives fewer than $k+1$ goods while another agent j receives more than $k+1$ goods, so agent i envies agent j by at least two goods, thereby violating EF1. Let $\mathcal{B} = (B_1, \dots, B_n)$ be the allocation such that agent 1 receives $k+2$ goods, agent 2 receives k goods, and every other agent receives $k+1$ goods. We have $\sum_{i \in N} f(u_i(B_i)) = f((k+2)a) + f(kb) + (n-2)f((k+1)b)$. Since \mathcal{A} is the only EF1 allocation, the additive welfarist rule with f chooses (only) \mathcal{A} , and it holds that $\sum_{i \in N} f(u_i(A_i)) > \sum_{i \in N} f(u_i(B_i))$. Rearranging the terms, we get

$$f((k+1)b) - f(kb) > f((k+2)a) - f((k+1)a),$$

or equivalently, $\Delta_{f,k}(b) > \Delta_{f,k+1}(a)$. Since k, a, b were arbitrarily chosen, $\Delta_{f,k}(b) > \Delta_{f,k+1}(a)$ holds for all $k \in \mathbb{Z}_{\geq 0}$ and $a, b \in \mathbb{R}_{> 0}$. Therefore, f satisfies Condition 1. \square

Next, we show that Condition 1 is sufficient for the function f to satisfy Theorem 3.1(b). To this end, we prove in Lemmas 3.3 and 3.4 that Condition 1 implies Condition 1a, which in turn implies that f satisfies the statement in Theorem 3.1(b). The proofs are rather complex and involve careful analyses of functions; we outline their main steps here and leave the details to our full version [8].

LEMMA 3.3. *Let $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R} \cup \{-\infty\}$ be a strictly increasing function that satisfies Condition 1. Then, f satisfies Condition 1a.*

PROOF SKETCH. Let $k \in \mathbb{Z}_{> 0}$ be given. We show that $\sup \Delta_{f,k}$ and $\inf \Delta_{f,k}$ are both finite, and hence $d_{f,k} := \sup \Delta_{f,k} - \inf \Delta_{f,k}$ is well-defined and non-negative. We then prove that $d_{f,k} = 0$, so $\Delta_{f,k}$ is a constant function, and f satisfies Condition 1a. \square

Next, we state another key lemma. Combined with Lemma 3.3, this lemma shows that if f satisfies Condition 1, then it satisfies the statement in Theorem 3.1(b).

LEMMA 3.4. *Let $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R} \cup \{-\infty\}$ be a strictly increasing function continuous on $\mathbb{R}_{> 0}$ that satisfies Condition 1a. Then, there exist constants $\alpha \in \mathbb{R}_{> 0}$ and $\beta \in \mathbb{R}$ such that $f(x) = \alpha \log x + \beta$ for all $x \in \mathbb{R}_{\geq 0}$.*

PROOF SKETCH. Define f_0 such that $f_0(x) = \alpha \log x + \beta$ where $\alpha = (f(2) - f(1))/\log 2 \in \mathbb{R}_{> 0}$ and $\beta = f(1) \in \mathbb{R}$. We show that f and f_0 coincide on the set $\{2^t : t \text{ is rational}\}$. Since this set is dense in $\mathbb{R}_{> 0}$ and f (and f_0) is continuous, f and f_0 coincide on $\mathbb{R}_{\geq 0}$. \square

We remark that Suksompong [21, Lemma 2] proved a result similar to Lemma 3.4 but made the stronger assumption that the function f is differentiable. Our result only assumes continuity.

We are now ready to prove Theorem 3.1.

PROOF OF THEOREM 3.1. The implication (a) \Rightarrow (b) follows from Lemmas 3.2, 3.3, and 3.4, while the implication (b) \Rightarrow (a) follows from the result of Caragiannis et al. [6] that MNW guarantees EF1. \square

Theorem 3.1 and Lemmas 3.2, 3.3, and 3.4 imply that a continuous function having the logarithmic property is equivalent to the function satisfying Conditions 1 and 1a (separately). We view this as an interesting mathematical property in itself.

3.2 Two-Value Instances

We now consider two-value instances, which are instances with only two possible utilities for the goods. Note that the classes of identical-good instances and two-value instances do not contain each other. Indeed, an identical-good instance with $n \geq 3$ agents may be n -value rather than two-value since each of the n agents could assign a unique utility to the goods,⁹ whereas a two-value instance allows an agent to have varying utilities for different goods.

Despite this incomparability, we show that the same characterization for identical-good instances also holds for two-value instances.

THEOREM 3.5. *Let $n \geq 2$ be given, and let $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R} \cup \{-\infty\}$ be a strictly increasing function continuous on $\mathbb{R}_{> 0}$. Then, the following statements are equivalent:*

- (a) *For every positive-admitting two-value instance with n agents, every allocation chosen by the additive welfarist rule with f is EF1.*
- (b) *There exist constants $\alpha \in \mathbb{R}_{> 0}$ and $\beta \in \mathbb{R}$ such that $f(x) = \alpha \log x + \beta$ for all $x \in \mathbb{R}_{\geq 0}$.*

3.3 Normalized Instances

Next, we consider normalized instances, where agents assign the same utility to the entire set of goods M . We first consider the case of three or more agents. We show that just like for the cases of identical-good instances and two-value instances, the only additive welfarist rule that yields EF1 for this case is the MNW rule. This extends Theorem 6 of Eckart et al. [10], which only handles p -mean rules (as opposed to all additive welfarist rules).

THEOREM 3.6. *Let $n \geq 3$ be given, and let $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R} \cup \{-\infty\}$ be a strictly increasing function continuous on $\mathbb{R}_{> 0}$. Then, the following statements are equivalent:*

- (a) *For every positive-admitting normalized instance with n agents, every allocation chosen by the additive welfarist rule with f is EF1.*
- (b) *There exist constants $\alpha \in \mathbb{R}_{> 0}$ and $\beta \in \mathbb{R}$ such that $f(x) = \alpha \log x + \beta$ for all $x \in \mathbb{R}_{\geq 0}$.*

To prove Theorem 3.6, we cannot use the construction in the proof of Lemma 3.2 since the instance in that construction is not necessarily normalized. Instead, we show that the function defining the additive welfarist rule must satisfy Condition 1a. To this end, we augment the construction of Suksompong [21] by adding a highly valuable good and an extra agent who only values that good, so

⁹However, for two agents, an identical-good instance is necessarily two-value.

that the instance becomes normalized. The full proof is deferred to the full version of this paper [8].

We now address the case of two agents. Eckart et al. [10] showed that in this case, the p -mean rule guarantees EF1 for all $p \leq 0$, which implies that there is a larger class of additive welfarist rules that guarantee EF1. Accordingly, our characterization in Theorem 3.6 does not work for two agents. The problem of finding a characterization for this case turns out to be rather challenging. We instead provide a necessary condition and a sufficient condition for the functions defining the additive welfarist rules that guarantee EF1 for normalized instances with two agents. Note that the sufficient condition in Proposition 3.8 is a generalization of the condition used by Eckart et al. [10, Lemma 3].

PROPOSITION 3.7. *Let $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R} \cup \{-\infty\}$ be a strictly increasing function continuous on $\mathbb{R}_{>0}$ such that for every positive-admitting normalized instance with two agents, every allocation chosen by the additive welfarist rule with f is EF1. Then, f is strictly concave.*

PROPOSITION 3.8. *Let $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R} \cup \{-\infty\}$ be a strictly increasing function that satisfies Condition 2. Then, for every positive-admitting normalized instance with two agents, every allocation chosen by the additive welfarist rule with f is EF1.*

We show in our full version [8] that beyond the p -mean rules for $p \leq 0$, there are (infinitely many) other additive welfarist rules that guarantee EF1 for normalized instances with two agents. In particular, any linear combination of φ_p —the function defining the p -mean rule—satisfies Condition 2, so the additive welfarist rule with such a function also guarantees EF1 for this class of instances. On the other hand, the additive welfarist rule with the modified logarithmic function λ_c where $c > 0$ and the additive welfarist rule with the modified harmonic function h_c where $c > -1$ do not guarantee EF1 for these instances.

4 INTEGER-VALUED INSTANCES

In this section, we turn our attention to *integer-valued* instances, where the utility of each agent for each good must be a (non-negative) integer. For these instances, the MNW rule is no longer the unique additive welfarist rule that guarantees EF1: the MHW rule exhibits the same property [17]. We shall explore the conditions for rules to satisfy this property, and provide several examples of such rules.

4.1 Identical-Good Instances

We begin with the class of (integer-valued) identical-good instances. We show that the functions defining the additive welfarist rules that guarantee EF1 for such instances are precisely those that satisfy Condition 3. Perhaps unsurprisingly, Condition 3 is similar to Condition 1 (which is the condition corresponding to *real-valued* identical-good instances), with the difference being that the values of a and b in the conditions are positive real numbers in Condition 1 and positive integers in Condition 3.

THEOREM 4.1. *Let $n \geq 2$ be given, and let $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R} \cup \{-\infty\}$ be a strictly increasing function. Then, the following statements are equivalent:*

- (a) *For every positive-admitting integer-valued identical-good instance with n agents, every allocation chosen by the additive welfarist rule with f is EF1.*
- (b) *f satisfies Condition 3.*

Due to the similarity between Conditions 1 and 3, the proof for (a) \Rightarrow (b) in Theorem 4.1 follows similarly to that in the real-valued case (Lemma 3.2). To prove (b) \Rightarrow (a), we show that Condition 3 implies Condition 3a, which in turn implies (a).

LEMMA 4.2. *Let $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R} \cup \{-\infty\}$ be a strictly increasing function that satisfies Condition 3. Then, f satisfies Condition 3a.*

PROOF. Let $k, \ell \in \mathbb{Z}_{\geq 0}$ and $a, b \in \mathbb{Z}_{>0}$ be given such that $\ell < k$. By applying Condition 3 repeatedly, we have $\Delta_{f,\ell}(b) > \Delta_{f,\ell+1}(a) > \dots > \Delta_{f,k}(a)$, so f indeed satisfies Condition 3a. \square

LEMMA 4.3. *Let $n \geq 2$ be given, and let $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R} \cup \{-\infty\}$ be a strictly increasing function that satisfies Condition 3a. Then, the statement in Theorem 4.1(a) holds.*

PROOF OF THEOREM 4.1. The implication (a) \Rightarrow (b) follows similarly as Lemma 3.2, while the implication (b) \Rightarrow (a) follows from Lemmas 4.2 and 4.3. \square

Since Conditions 1 and 1a are equivalent (see Theorem 3.1 and Lemmas 3.2, 3.3, and 3.4), one may be tempted to think that there is a condition analogous to Condition 1a that Condition 3 is equivalent to, perhaps “ $\Delta_{f,k}$ is a constant function (on domain $\mathbb{Z}_{>0}$) for each $k \in \mathbb{Z}_{>0}$ ”. However, this condition is too strong, and is not implied by Condition 3.¹⁰ Instead, we show that Condition 3 is equivalent to Condition 3b.

PROPOSITION 4.4. *Let $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R} \cup \{-\infty\}$ be a strictly increasing function. Then, the following statements are equivalent:*

- (a) *f satisfies Condition 3.*
- (b) *f satisfies Condition 3a.*
- (c) *f satisfies Condition 3b.*

We next give examples of additive welfarist rules that guarantee EF1 for all integer-valued identical-good instances. We study two families of functions: modified logarithmic functions λ_c and modified harmonic numbers h_c . By leveraging Proposition 4.4 and Theorem 4.1, we can determine which of these additive welfarist rules ensure EF1 for integer-valued identical-good instances.

PROPOSITION 4.5. *The function λ_c satisfies Condition 3b if and only if $0 \leq c \leq 1$.*

PROPOSITION 4.6. *The function h_c satisfies Condition 3b if and only if $-1 \leq c \leq 1/\log 2 - 1$ (≈ 0.443).*

Propositions 4.5 and 4.6 show that beyond the MNW rule and the MHW rule, many other additive welfarist rules guarantee EF1 for integer-valued identical-good instances. We provide more discussion in the full version of our paper [8].

¹⁰Indeed, the MHW rule—which corresponds to the function h_0 —guarantees EF1 for integer-valued instances [17], but $\Delta_{h_0,1}$ is not a constant function as $\Delta_{h_0,1}(1) = h_0(2) - h_0(1) = 1/2 < 7/12 = h_0(4) - h_0(2) = \Delta_{h_0,1}(2)$.

4.2 Binary Instances

Next, we consider binary instances, where each good is worth either 0 or 1 to each agent. We show that the functions defining the additive welfarist rules that guarantee EF1 for all binary instances are those that satisfy *Condition 4*, which is equivalent to the function being strictly concave in the domain of non-negative integers.

THEOREM 4.7. *Let $n \geq 2$ be given, and let $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R} \cup \{-\infty\}$ be a strictly increasing function. Then, the following statements are equivalent:*

- (a) *For every positive-admitting binary instance with n agents, every allocation chosen by the additive welfarist rule with f is EF1.*
- (b) *f satisfies Condition 4.*

PROOF. (a) \Rightarrow (b): Let $k \in \mathbb{Z}_{\geq 0}$. Consider an instance with n agents and $2k + n$ goods, and let $M' = \{g_1, \dots, g_{2k+2}\}$. The utilities of the goods are as follows.

- For $i \in \{1, 2\}$, $u_i(g) = 1$ for $g \in M'$, and $u_i(g) = 0$ otherwise.
- For $i \in \{3, \dots, n\}$, $u_i(g_{2k+i}) = 1$, and $u_i(g) = 0$ for all other $g \in M'$.

This instance is a binary instance; moreover, it is positive-admitting since any allocation where each agent $i \in N$ receives g_{2k+i} gives positive utility to every agent.

Let $\mathcal{A} = (A_1, \dots, A_n)$ be an allocation chosen by the additive welfarist rule with f . Each good must be allocated to an agent who has utility 1 for the good; otherwise, transferring such a good to an agent who has utility 1 for the good increases the value of $\sum_{i \in N} f(u_i(A_i))$, contradicting the assumption that \mathcal{A} is chosen by the additive welfarist rule with f . Accordingly, the goods in M' are allocated to agent 1 and 2, and for each $i \in \{3, \dots, n\}$, g_{2k+i} is allocated to agent i .

Recall that \mathcal{A} is EF1 since it is chosen by the additive welfarist rule with f . We claim that agent 1 and 2 each receives exactly $k + 1$ goods from M' . If not, then one of them receives at most k goods from M' , and will envy the other agent (who receives at least $k + 2$ goods from M') by more than one good, making the allocation not EF1 and contradicting our assumption. Then, we have $\sum_{i \in N} f(u_i(A_i)) = 2f(k + 1) + (n - 2)f(1)$.

Let $\mathcal{B} = (B_1, \dots, B_n)$ be the allocation such that agent 1 receives k goods from M' , agent 2 receives $k + 2$ goods from M' , and for each $i \in \{3, \dots, n\}$, agent i receives g_{2k+i} . We have $\sum_{i \in N} f(u_i(B_i)) = f(k) + f(k + 2) + (n - 2)f(1)$. Note that \mathcal{B} is not EF1, and cannot be chosen by the additive welfarist rule with f . Therefore, we have $\sum_{i \in N} f(u_i(A_i)) > \sum_{i \in N} f(u_i(B_i))$. Rearranging the terms, we get

$$f(k + 1) - f(k) > f(k + 2) - f(k + 1),$$

or equivalently, $\Delta_{f,k}(1) > \Delta_{f,k+1}(1)$. Since $k \in \mathbb{Z}_{\geq 0}$ was arbitrarily chosen, $\Delta_{f,k}(1) > \Delta_{f,k+1}(1)$ holds for all $k \in \mathbb{Z}_{\geq 0}$. Therefore, f satisfies Condition 4.

(b) \Rightarrow (a): Let a positive-admitting binary instance with n agents be given, and let $\mathcal{A} = (A_1, \dots, A_n)$ be an allocation chosen by the additive welfarist rule with f . Assume, for the sake of contradiction, that \mathcal{A} is not EF1. Then, there exist $i, j \in N$ such that agent i envies agent j by more than one good, i.e., $u_i(A_i) < u_i(A_j \setminus \{g\})$ for all $g \in A_j$. Note that we must have $u_i(g) \leq u_j(g)$ for all $g \in A_j$. Indeed, otherwise we have $u_i(g) = 1$ and $u_j(g) = 0$, and transferring g from

agent j 's bundle to agent i 's bundle increases $f(u_i(A_i))$ and does not decrease $f(u_j(A_j))$, thereby increasing $\sum_{k \in N} f(u_k(A_k))$ and contradicting the assumption that \mathcal{A} is chosen by the additive welfarist rule with f . Moreover, since $u_i(A_j) > 0$, there exists $g' \in A_j$ such that $u_i(g') = 1$ (and hence $u_j(g') = 1$).

Let $\mathcal{B} = (B_1, \dots, B_n)$ be the same allocation as \mathcal{A} except that g' is transferred from agent j 's bundle to agent i 's bundle, i.e., $B_i = A_i \cup \{g'\}$, $B_j = A_j \setminus \{g'\}$, and $B_k = A_k$ for all $k \in N \setminus \{i, j\}$. Note that $u_i(A_i) < u_i(A_j \setminus \{g'\}) = u_i(B_j)$, and that $u_i(B_j) \leq u_j(B_j)$ since $u_i(g) \leq u_j(g)$ for all $g \in B_j$. Therefore, $u_i(A_i) < u_j(B_j)$. By Condition 4, it holds that $\Delta_{f,u_i(A_i)}(1) > \dots > \Delta_{f,u_j(B_j)}(1)$. On the other hand, we have $\sum_{k \in N} f(u_k(B_k)) = f(u_i(B_i)) + f(u_j(B_j)) + \sum_{k \in N \setminus \{i, j\}} f(u_k(A_k))$. Since \mathcal{A} is chosen by the additive welfarist rule with f , we have $\sum_{k \in N} f(u_k(B_k)) \leq \sum_{k \in N} f(u_k(A_k))$. Rearranging the terms, we get

$$f(u_i(B_i)) - f(u_i(A_i)) \leq f(u_j(A_j)) - f(u_j(B_j)),$$

or equivalently, $\Delta_{f,u_i(A_i)}(1) \leq \Delta_{f,u_j(B_j)}(1)$, a contradiction. Therefore, \mathcal{A} is EF1. \square

Condition 4 is weaker than Condition 3. Indeed, if f satisfies Condition 3, then it satisfies Condition 4 by definition. However, the converse is not true: the function $f(x) = \sqrt{x}$ satisfies Condition 4, but it does not satisfy Condition 3 since $\Delta_{f,0}(1) = 1$ is not greater than $\Delta_{f,1}(6) \approx 1.01$. This means that each additive welfarist rule that guarantees EF1 for every positive-admitting integer-valued identical-good instance also guarantees EF1 for every positive-admitting binary instance, but the converse does not hold.

PROPOSITION 4.8. *Let $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R} \cup \{-\infty\}$ be a strictly increasing function. If f satisfies Condition 3, then it satisfies Condition 4.*

We mentioned earlier that Condition 4 is equivalent to the function being strictly concave in the domain of non-negative integers. On the other hand, the function φ_p defining the p -mean rule is strictly concave if and only if $p < 1$. We prove that, indeed, φ_p satisfies Condition 4 exactly when $p < 1$.

PROPOSITION 4.9. *The function φ_p satisfies Condition 4 if and only if $p < 1$.*

4.3 Two-Value Instances

We now consider integer-valued instances which are two-value, a strict generalization of binary instances. Recall from Section 3.2 that the classes of identical-good instances and two-value instances are not subclasses of each other (except when $n = 2$). Therefore, the techniques used in Section 4.1 cannot be used in this section.¹¹

We show that the characterization for integer-valued two-value instances is given by *Condition 5*. Before we state the characterization, we examine the relationship between Conditions 3 and 5.

PROPOSITION 4.10. *Let $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R} \cup \{-\infty\}$ be a strictly increasing function.*

- (i) *If f satisfies Condition 5, then f satisfies Condition 3.*
- (ii) *If f satisfies Condition 3 and the property that for each $k \in \mathbb{Z}_{> 0}$, the function $\Delta_{f,k}$ is either non-decreasing or non-increasing on the domain of positive integers, then f satisfies Condition 5.*

¹¹In particular, the proof that Condition 3a is a sufficient condition for the identical-good case (Lemma 4.3) does not apply to the two-value case.

We now state the characterization for integer-valued two-value instances.

THEOREM 4.11. *Let $n \geq 2$ be given, and let $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R} \cup \{-\infty\}$ be a strictly increasing function. Then, the following statements are equivalent:*

- (a) *For every positive-admitting integer-valued two-value instance with n agents, every allocation chosen by the additive welfarist rule with f is EF1.*
- (b) *f satisfies Condition 5.*

Proposition 4.10 says that Condition 5 is stronger than Condition 3. Therefore, Theorem 4.11 implies that every additive welfarist rule that guarantees EF1 for every positive-admitting integer-valued two-value instance also guarantees EF1 for every positive-admitting integer-valued identical-good instance, even though these two classes of instances are not subclasses of each other.

Next, we give examples of additive welfarist rules that ensure EF1 for all integer-valued two-value instances. We use Propositions 4.5, 4.6, and 4.10 to prove these results.

PROPOSITION 4.12. *The function λ_c satisfies Condition 5 if and only if $0 \leq c \leq 1$.*

PROPOSITION 4.13. *The function h_c satisfies Condition 5 if and only if $-1 \leq c \leq 1/\log 2 - 1$ (≈ 0.443).*

4.4 General Integer-Valued Instances

Finally, we consider the class of all integer-valued instances. It turns out to be challenging to find an exact characterization for this class of instances. Instead, we provide a necessary condition in the form of Condition 6a and a sufficient condition in the form of Condition 6b.

PROPOSITION 4.14. *Let $n \geq 2$ be given, and let $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R} \cup \{-\infty\}$ be a strictly increasing function such that for every positive-admitting integer-valued instance with n agents, every allocation chosen by the additive welfarist rule with f is EF1. Then, f satisfies Condition 6a.*

PROPOSITION 4.15. *Let $n \geq 2$ be given, and let $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R} \cup \{-\infty\}$ be a strictly increasing function that satisfies Condition 6b. Then, for every positive-admitting integer-valued instance with n agents, every allocation chosen by the additive welfarist rule with f is EF1.*

We demonstrate that the necessary condition given as Condition 6a is fairly tight, in the sense that it is stronger than Condition 5, which characterizes the functions defining the additive welfarist rules that guarantee EF1 for integer-valued two-value instances.

PROPOSITION 4.16. *Let $f : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R} \cup \{-\infty\}$ be a strictly increasing function. If f satisfies Condition 6a, then it satisfies Condition 5.*

We now present some examples. We show that just like for the subclasses of integer-valued identical-good instances (Proposition 4.5), binary instances (Proposition 4.8), and integer-valued two-value instances (Proposition 4.12), the additive welfarist rule with function λ_c for $0 \leq c \leq 1$ guarantees EF1 for all integer-valued instances.

PROPOSITION 4.17. *The function λ_c satisfies Condition 6b if and only if $0 \leq c \leq 1$.*

On the other hand, leveraging Condition 6a, we show that the additive welfarist rule with function h_c for $-1 \leq c < -1/2$ does not guarantee EF1 for the class of all integer-valued instances. This stands in contrast to integer-valued identical-good instances (Proposition 4.6), binary instances (Proposition 4.8), and integer-valued two-value instances (Proposition 4.13).

PROPOSITION 4.18. *The function h_c does not satisfy Condition 6a when $-1 \leq c < -1/2$.*

5 CONCLUSION

In this paper, we have characterized additive welfarist rules that guarantee EF1 allocations for various classes of instances. In the real-valued case, we strengthened the result of Suksompong [21] by showing that only the maximum Nash welfare (MNW) rule ensures EF1 even for the most restricted class of identical-good instances. On the other hand, in the practically important case where all values are integers, we demonstrated that even for the most general setting with no additional restrictions, there is a wide range of additive welfarist rules that always return EF1 allocations.

Since we have established the existence of several alternatives that perform as well as MNW in terms of guaranteeing EF1 allocations for integer-valued instances, a natural follow-up would be to compare these rules with respect to other measures, both to MNW and to one another. As illustrated in Example 1.1, some additive welfarist rules may produce more preferable allocations for certain instances than others. It would be interesting to formalize this observation, for example, by comparing their prices in terms of social welfare, in the same vein as the “price of fairness” [3, 7, 15].

Other possible future directions include completely characterizing additive welfarist rules for normalized instances with two agents (Section 3.3) and for general integer-valued instances (Section 4.4) by tightening their respective necessary and sufficient conditions. Obtaining characterizations of welfarist rules that are not necessarily additive is also a possible extension of our work. Beyond EF1, one could consider other fairness notions from the literature, such as proportionality up to one good (PROP1). Finally, another intriguing avenue is to study the problem of computing (or approximating) an allocation produced by various additive welfarist rules, as has been done extensively for the MNW rule [1, 2, 9, 14].

ACKNOWLEDGMENTS

This work was partially supported by the Singapore Ministry of Education under grant number MOE-T2EP20221-0001 and by an NUS Start-up Grant. We thank the anonymous reviewers of AAMAS 2025 for their valuable comments, as well as Sanjay Jain and Frank Stephan for their help with part of the proof of Proposition 4.6.

REFERENCES

- [1] Hannaneh Akrami, Bhaskar Ray Chaudhury, Martin Hoefer, Kurt Mehlhorn, Marco Schmalhofer, Golnoosh Shahkarami, Giovanna Varricchio, Quentin Vermande, and Ernest van Wijland. 2022. Maximizing Nash social welfare in 2-value instances. In *Proceedings of the 36th AAAI Conference on Artificial Intelligence (AAAI)*. 4760–4767.
- [2] Georgios Amanatidis, Georgios Birmpas, Aris Filos-Ratsikas, Alexandros Hollender, and Alexandros A. Voudouris. 2021. Maximum Nash welfare and other stories about EFX. *Theoretical Computer Science* 863 (2021), 69–85.

- [3] Xiaohui Bei, Xinhang Lu, Pasin Manurangsi, and Warut Suksompong. 2021. The price of fairness for indivisible goods. *Theory of Computing Systems* 65, 7 (2021), 1069–1093.
- [4] Steven J. Brams and Alan D. Taylor. 1996. *Fair Division: From Cake-Cutting to Dispute Resolution*. Cambridge University Press.
- [5] Eric Budish. 2011. The combinatorial assignment problem: Approximate competitive equilibrium from equal incomes. *Journal of Political Economy* 119, 6 (2011), 1061–1103.
- [6] Ioannis Caragiannis, David Kurokawa, Hervé Moulin, Ariel D. Procaccia, Nisarg Shah, and Junxing Wang. 2019. The unreasonable fairness of maximum Nash welfare. *ACM Transactions on Economics and Computation* 7, 3 (2019), 12:1–12:32.
- [7] Karen Frilya Celine, Muhammad Ayaz Dzulfikar, and Ivan Adrian Koswara. 2023. Egalitarian price of fairness for indivisible goods. In *Proceedings of the 20th Pacific Rim International Conference on Artificial Intelligence (PRICAI)*, 23–28.
- [8] Karen Frilya Celine, Warut Suksompong, and Sheung Man Yuen. 2024. On the fairness of additive welfarist rules. *arXiv preprint arXiv:2412.15472* (2024).
- [9] Richard Cole and Vasilis Gkatzelis. 2018. Approximating the Nash social welfare with indivisible items. *SIAM J. Comput.* 47, 3 (2018), 1211–1236.
- [10] Owen Eckart, Alexandros Psomas, and Paritosh Verma. 2024. On the fairness of normalized p -means for allocating goods and chores. In *Proceedings of the 25th ACM Conference on Economics and Computation (EC)*, 1267. Extended version available at arXiv:2402.14996v1.
- [11] Duncan K. Foley. 1967. Resource allocation and the public sector. *Yale Economics Essays* 7, 1 (1967), 45–98.
- [12] Jonathan Goldman and Ariel D. Procaccia. 2014. Spliddit: Unleashing fair division algorithms. *ACM SIGecom Exchanges* 13, 2 (2014), 41–46.
- [13] Wolfgang Hintze. 2019. Analytic continuation of harmonic series. Mathematics Stack Exchange. <https://math.stackexchange.com/q/3058569> Accessed 30-09-2024.
- [14] Euiwoong Lee. 2017. APX-hardness of maximizing Nash social welfare with indivisible items. *Inform. Process. Lett.* 122 (2017), 17–20.
- [15] Zihao Li, Shengxin Liu, Xinhang Lu, Biaoshuai Tao, and Yichen Tao. 2024. A complete landscape for the price of envy-freeness. In *Proceedings of the 23rd International Conference on Autonomous Agents and Multiagent Systems (AAMAS)*, 1183–1191.
- [16] Richard J. Lipton, Evangelos Markakis, Elchanan Mossel, and Amin Saberi. 2004. On approximately fair allocations of indivisible goods. In *Proceedings of the 5th ACM Conference on Electronic Commerce (EC)*, 125–131.
- [17] Luisa Montanari, Ulrike Schmidt-Kraepelin, Warut Suksompong, and Nicholas Teh. 2025. Weighted envy-freeness for submodular valuations. *Social Choice and Welfare* 65, 4 (2025), 983–1016.
- [18] Hervé Moulin. 2003. *Fair Division and Collective Welfare*. MIT Press.
- [19] Hervé Moulin. 2019. Fair division in the Internet age. *Annual Review of Economics* 11 (2019), 407–441.
- [20] Jack Robertson and William Webb. 1998. *Cake-Cutting Algorithms: Be Fair if You Can*. Peters/CRC Press.
- [21] Warut Suksompong. 2023. A characterization of maximum Nash welfare for indivisible goods. *Economics Letters* 222 (2023), 110956.
- [22] William Thomson. 2016. Introduction to the theory of fair allocation. In *Handbook of Computational Social Choice*, Felix Brandt, Vincent Conitzer, Ulle Endriss, Jérôme Lang, and Ariel D. Procaccia (Eds.). Cambridge University Press, Chapter 11, 261–283.
- [23] Hal R. Varian. 1974. Equity, envy, and efficiency. *Journal of Economic Theory* 9 (1974), 63–91.
- [24] Sheung Man Yuen and Warut Suksompong. 2023. Extending the characterization of maximum Nash welfare. *Economics Letters* 224 (2023), 111030.